## Pending Claims Application No. 09/685,577

<u>;</u>

Attorney Docket No.: 05725.0656-01000 Filed: October 11, 2000

 (Previously presented) A structured cosmetic composition comprising:

at least one continuous liquid fatty phase,

wherein said at least one continuous liquid fatty phase is structured with a sufficient amount of at least one polymer of formula (I) and mixtures thereof:

$$R^{1} \longrightarrow O = \begin{cases} C \longrightarrow R^{2} \longrightarrow C \longrightarrow N \longrightarrow R^{3} \longrightarrow N \longrightarrow C \longrightarrow R^{2} \longrightarrow C \longrightarrow C \longrightarrow R^{1} \\ 0 \longrightarrow O \longrightarrow N \longrightarrow C \longrightarrow R^{2} \longrightarrow C \longrightarrow C \longrightarrow R^{1} \\ 0 \longrightarrow O \longrightarrow N \longrightarrow C \longrightarrow C \longrightarrow R^{2} \longrightarrow C \longrightarrow R^$$

in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;
- R<sup>1</sup>, which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;
- $R^2$ , which are identical or different, are each chosen from  $C_4$  to  $C_{42}$  hydrocarbon-based groups with the proviso that at least 50% of  $R^2$  are chosen from  $C_{30}$  to  $C_{42}$  hydrocarbon-based groups;

- R<sup>3</sup>, which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms and nitrogen atoms with the proviso that R<sup>3</sup> comprises at least 2 carbon atoms; and

بج

- R<sup>4</sup>, which are identical or different, are each chosen from hydrogen atoms, C<sub>1</sub> to C<sub>10</sub> alkyl groups and a direct bond to group chosen from R<sup>3</sup> and another R<sup>4</sup> such that when said at least one group is chosen from another R<sup>4</sup>, the nitrogen atom to which both R<sup>3</sup> and R<sup>4</sup> are bonded forms part of a heterocyclic structure defined in part by R<sup>4</sup>-N-R<sup>3</sup>, with the proviso that at least 50% of all R<sup>4</sup> are chosen from hydrogen atoms;

wherein said structured composition is in the form of a non-migrating, waxfree solid, and

wherein said at least one continuous liquid fatty phase and said at least one polymer form a physiologically acceptable medium.

- 2 46. (Canceled)
- 47. (Original) A composition according to Claim 1, further comprising at least one amphiphilic compound chosen from amphiphilic compounds which are liquid at room temperature and have an HLB value of less than 12.
- 48. (Original) A composition according to Claim 47, wherein said HLB value ranges from 1 to 7.
- 49. (Original) A composition according to Claim 47, wherein said HLB value ranges from 1 to 5.
  - 50 149. (Canceled)

150. (Previously presented) A process of structuring a cosmetic composition in the form of a physiologically acceptable composition, which is wax-free and non-migrating comprising including in said composition at least one liquid continuous fatty phase, said at least one liquid continuous fatty phase being structured with a sufficient amount of at least one polymer of formula (I) and mixtures thereof:

in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;
- R<sup>1</sup>, which are identical or different, are each chosen from alkyl groups comprising at least 4 carbon atoms and alkenyl groups comprising at least 4 carbon atoms;
- $R^2$ , which are identical or different, are each chosen from  $C_4$  to  $C_{42}$  hydrocarbon-based groups with the proviso that at least 50% of  $R^2$  are chosen from  $C_{30}$  to  $C_{42}$  hydrocarbon-based groups;
- R<sup>3</sup>, which are identical or different, are each chosen from organic groups comprising atoms chosen from carbon atoms, hydrogen atoms, oxygen atoms

ت

and nitrogen atoms with the proviso that R<sup>3</sup> comprises at least 2 carbon atoms; and

- R<sup>4</sup>, which are identical or different, are each chosen from hydrogen atoms, C<sub>1</sub> to C<sub>10</sub> alkyl groups and a direct bond to group chosen from R<sup>3</sup> and another R<sup>4</sup> such that when said at least one group is chosen from another R<sup>4</sup>, the nitrogen atom to which both R<sup>3</sup> and R<sup>4</sup> are bonded forms part of a heterocyclic structure defined in part by R<sup>4</sup>-N-R<sup>3</sup>, with the proviso that at least 50% of all R<sup>4</sup> are chosen from hydrogen atoms;and

wherein said composition is wax-free and non-migrating.

- 151 153. (Canceled)
- 154. (Original) A process according to Claim 150, wherein said at least one structuring polymer is combined with at least one amphiphilic compound that is liquid at room temperature, with an HLB value of less than 12.
- 155. (Original) A process according to Claim 154, wherein said HLB ranges from 1 to 7.
- 156. (Original) A process according to Claim 155, wherein said HLB ranges from 1 to 5.

157 - 188. (Canceled)